

1 Lie bracket

Recall that vector fields can be identified with homogeneous linear first order partial differential operators $X = X^a \partial_a$ acting on functions as $X(f) = X^a \partial_a f$.

The Lie-bracket $[X, Y]$ of two vector fields X and Y is defined as

$$[X, Y](f) = X(Y(f)) - Y(X(f)) .$$

Show that $[X, Y]$ also is a vector field, i.e. a homogeneous linear first order differential operator, with components

$$[X, Y]^a = X^b \partial_b Y^a - Y^b \partial_b X^a . \quad (1)$$

Check, by a direct coordinate calculation, that the right-hand-side of (1) transforms correctly under changes of coordinates.

Prove the *Jacobi identity*:

$$[X, [Y, Z]] + [Y, [Z, X]] + [Z, [X, Y]] = 0 .$$

2 Christoffel symbols

The Christoffel symbols of the Levi-Civita connection are defined by the formula

$$\Gamma_{bc}^a = \frac{1}{2} g^{ad} (\partial_b g_{cd} + \partial_c g_{bd} - \partial_d g_{bc}) .$$

Using this definition, calculate the Christoffel symbols for a) the Euclidean metric on \mathbb{R}^2 in polar coordinates: $g = d\rho^2 + \rho^2 d\varphi^2$, and b) the unit round metric on S^2 : $h = d\theta^2 + \sin^2 \theta d\varphi^2$.

Show that the Euler-Lagrange equations

$$\frac{d}{dt} \left(\frac{\partial L}{\partial \dot{x}^a} \right) = \frac{\partial L}{\partial x^a} \quad (2)$$

associated with the Lagrange function

$$L(x^c, \dot{x}^c) = \frac{1}{2} g_{ab} \dot{x}^a \dot{x}^b \quad (3)$$

can be written as

$$\ddot{x}^a + \Gamma_{bc}^a \dot{x}^b \dot{x}^c = 0 . \quad (4)$$

Use the explicit form of (2) for the two-dimensional metrics g and h above to calculate again the Christoffel symbols.

3 Geodesics

Solutions $x^a(t)$ of (4) are called *affinely parameterized geodesics*, and the parameter t is said to be *affine*.

Let $x^a(\sigma)$ be a geodesic with an affine parameter σ and let $W^a = dx^a/d\sigma$. Show that:

$$\frac{d}{d\sigma}(g_{ab}W^aW^b) = 0.$$

Recall that a vector X is called *timelike* if $g(X, X) > 0$, *null* if $X \neq 0$ and $g(X, X) = 0$, and *spacelike* if $g(X, X) < 0$. Deduce that the type of the tangent vector does not change along a geodesic. Show that for a timelike geodesic, σ is an affine function of proper time τ (this means that $\sigma = a\tau + b$, for some constants a, b).