

- 1 Let a bracket over indices denote complete antisymmetrisation, and a parenthesis over indices denote complete symmetrisation: for example,

$$A_{[\mu\nu]} = \frac{1}{2}(A_{\mu\nu} - A_{\nu\mu}), \quad A_{(\mu\nu)} = \frac{1}{2}(A_{\mu\nu} + A_{\nu\mu}), \quad \delta_{\mu}^{[\alpha} \delta_{\rho}^{\gamma]} = \frac{1}{2}(\delta_{\mu}^{\alpha} \delta_{\rho}^{\gamma} - \delta_{\mu}^{\gamma} \delta_{\rho}^{\alpha}),$$

and similarly for 3, 4 or more indices. Show that

- i.  $A^{[\mu\nu]} B_{\mu\nu} = A^{\mu\nu} B_{[\mu\nu]}$ ,
- ii.  $A^{[\mu\nu\rho]} B_{\mu\nu\rho} = A^{\mu\nu\rho} B_{[\mu\nu\rho]}$ ,
- iii.  $\delta_{\mu}^{[\alpha} \delta_{\rho}^{\gamma]} = \delta_{[\mu}^{\alpha} \delta_{\rho]}^{\gamma}$ ,
- iv.  $\delta_{\mu}^{[\alpha} \delta_{\nu}^{\beta} \delta_{\rho}^{\gamma]} = \delta_{[\mu}^{\alpha} \delta_{\nu}^{\beta} \delta_{\rho]}^{\gamma}$ ,
- v.  $\epsilon^{\alpha\beta\gamma\delta} \epsilon_{\alpha\beta\gamma\rho} = -6\delta_{\rho}^{\delta}$ ,
- vi.  $\epsilon^{\alpha\beta\gamma\delta} \epsilon_{\alpha\beta\nu\rho} = -4\delta_{[\nu}^{\gamma} \delta_{\rho]}^{\delta}$ .

- 2 Assuming the tensorial transformation law of  $F^{\mu\nu}$ , derive the explicit formulae for the transformation laws of the electric and magnetic fields under a boost along the first coordinate axis.
- 3 Show that the contraction  $F_{\alpha\beta} F^{\alpha\beta}$  is invariant (more precisely, behaves as a scalar) under Lorentz transformations, while  $F^{\alpha\beta} *F_{\alpha\beta}$  either remains invariant, or changes sign. Express those contractions in terms of  $\vec{E}$  and  $\vec{B}$ .
- 4 Let  $\vec{E} \cdot \vec{B} = 0$ , and suppose that  $|\vec{E}|^2 \neq |\vec{B}|^2$ . Show that there exists a Lorentz frame in which either  $\vec{E}$  or  $\vec{B}$  vanishes. [Hint: apply a boost with  $\vec{v}$  proportional to  $\vec{E} \times \vec{B}$ .]
- 5 The *alternating tensor*  $\epsilon_{\alpha\beta\gamma\delta}$  is defined by the requirement that it changes sign under the permutation of any two indices (such tensors are called *totally antisymmetric*), and

$$\epsilon_{0123} = 1.$$

Does this indeed define  $\epsilon_{\alpha\beta\gamma\delta}$  uniquely? [Hint: What is the value of  $\epsilon_{\alpha\beta\gamma\delta}$  when some indices coincide?]

Define  $\epsilon^{\alpha\beta\gamma\delta}$  by raising the indices using *some* symmetric two-contravariant tensor  $\eta^{\mu\nu}$ , with inverse tensor  $\eta_{\mu\nu}$ , possibly, but not necessarily, equal to the Minkowski metric:

$$\epsilon^{\alpha\beta\gamma\delta} = \eta^{\alpha\mu} \eta^{\beta\nu} \eta^{\gamma\rho} \eta^{\delta\sigma} \epsilon_{\mu\nu\rho\sigma}.$$

Show that  $\epsilon^{\alpha\beta\gamma\delta}$  is totally antisymmetric and that

$$\epsilon^{0123} = \det \eta^{\alpha\beta}.$$

Similarly show that

$$\Lambda^{\alpha'}_{\alpha} \Lambda^{\beta'}_{\beta} \Lambda^{\gamma'}_{\gamma} \Lambda^{\delta'}_{\delta} \epsilon_{\alpha'\beta'\gamma'\delta'} = \det \Lambda \epsilon_{\alpha\beta\gamma\delta}.$$

How can this be generalised to other dimensions? or to the Euclidean metric? How many totally antisymmetric tensors with five indices are there in dimension  $n$ ,  $1 \leq n \leq 7$ ?

6 [For self-study, will not be done in class]

a) Assuming that  $\Lambda^\alpha_\beta$  is a Lorentz matrix, show that the matrix  $A^\alpha_\beta := \eta^{\alpha\beta} \Lambda^\mu_\beta \eta_{\mu\nu}$  is inverse to  $\Lambda^\alpha_\beta$ .

b) Recall that we required that  $F^{\mu\nu}$  transforms as a *two-contravariant tensor* under Lorentz transformations: if  $\bar{x}^\alpha = \Lambda^\alpha_\beta x^\beta + a^\alpha$ , then

$$\bar{F}^{\mu\nu}(\bar{x}) = \Lambda^\mu_\alpha \Lambda^\nu_\beta F^{\alpha\beta}(x),$$

and that  $F_{\mu\nu}$  has been defined as

$$F_{\mu\nu} := \eta_{\mu\alpha} \eta_{\nu\beta} F^{\alpha\beta}.$$

Use 1) to show that  $F_{\alpha\beta}$  transforms as

$$\bar{F}_{\mu\nu}(\bar{x}) = (\Lambda^{-1})^\alpha_\mu (\Lambda^{-1})^\beta_\nu F_{\alpha\beta}(x)$$

(this is called the *transformation law of a two-covariant tensor*).